


# Ten Reasons for Laparoendoscopic Repair of Hiatal Hernia: Case Presentation With Long-Term Follow-Up

The American Surgeon  
00(0) 1–3  
© The Author(s) 2020  
Article reuse guidelines:  
sagepub.com/journals-permissions  
DOI: 10.1177/0003134820933608  
journals.sagepub.com/home/asu  


Medhat Fanous, MD, FACS<sup>1</sup>

## Abstract

Laparoendoscopic hiatal hernia repair (LEHHR) involves laparoscopic repair of hiatal hernia with concomitant transoral incisionless fundoplication (TIF). The objective of this case presentation is to highlight the benefits of LEHHR in a patient with long term follow up. This patient is a 56-year-old woman with symptoms of gastroesophageal reflux disease for 40 years. Esophagogastroduodenoscopy (EGD) showed a 2 cm hiatal hernia. DeMeester score was 21.3. She underwent LEHHR 33 months ago. The patient underwent laparoscopic cholecystectomy for symptomatic biliary dyskinesia. This provided the opportunity to examine the operative anatomy. There were minimal adhesions to the liver. The partial fundoplication was intact. The angle of His was preserved. The fundus was spared from any adhesions as TIF utilizes the cardia rather than the fundus to create the wrap. The plane behind the stomach was undisturbed. LEHHR has 10 main benefits. Anatomical benefits result from the preservation of the angle of His. Functional benefits relate to a partial fundoplication which normalizes pH values. LEHHR avoids bleeding from short gastric vessels and the creation of a wrap when anatomical obstacles present. Strategic benefits are directed toward any subsequent revisional reflux surgery. The lack of adhesions, easy access to the base of left crus, and sparing the fundus render revisional surgery straightforward.

## Keywords

gastroesophageal reflux disease, hiatal hernia repair, laparoendoscopic repair of hiatal hernia, transoral incisionless fundoplication

## Introduction

Laparoendoscopic hiatal hernia repair (LEHHR) involves laparoscopic hiatal hernia repair (LHHR) with concomitant transoral incisionless fundoplication (TIF). This technique was first reported by Dr Ihde in 2011.<sup>1</sup> The aim of this case report is to highlight the advantages of LEHHR and demonstrate the operative findings noted during a subsequent laparoscopic procedure almost 3 years later.

## Methods

This case involves a 56-year-old woman with a medical history of osteoporosis, asthma, aspiration pneumonia, and depression. Physical examination revealed a body mass index of 21.7 Kg/m<sup>2</sup> and an extremely small torso. She had symptoms of gastroesophageal reflux disease (GERD) and laryngopharyngeal reflux for 40 years. She presented with heartburn, regurgitation, asthma-like symptoms, hoarseness,

and globus sensation. She used proton pump inhibitors (PPIs) for 10 years, including esomeprazole 40 mg for 7 years and rabeprazole 20 mg daily for 3 years, with poor control of her symptoms. Her GERD Health-Related Quality of Life (GERD-HRQL) Questionnaire, Reflux Symptoms Index (RSI) Questionnaire and GERD Symptom Score (GERSS) Questionnaire on PPI were 56, 42, and 44, respectively.


Diagnostic workup included upper gastrointestinal series which showed normal motility. Esophagogastroduodenoscopy (EGD) showed a 2 cm hiatal hernia and Hill's deformity II (Figure 1B). The

<sup>1</sup>Department of Surgery, Aspirus Iron River Hospital & Clinics, Iron River, MI, USA

### Corresponding Author:

Medhat Fanous, MD, FACS, Department of Surgery, Aspirus Iron River Hospital & Clinics, 1400 W Ice Lake Road, Iron River, MI 49935, USA.


Email: medhat.fanous@gmail.com



**Figure 1.** Retroflexed endoscopic views. (A) depiction of the cardial wrap by the Esophyx device; (B) preoperative view; (C) intraoperative view; and (D) postoperative view at 33 months.

wireless pH study for 48 hours yielded a DeMeester score of 21.3.

LEHHR was performed using 4 ports due to her extremely small torso (camera, liver retractor, and 2 working ports). Pars Lucida of the lesser omentum was divided and dissection continued along the edge of the right crus. The pharyngoesophageal membrane was divided. The adhesions to the left crus were divided. A retroesophageal window was made. The stomach was retracted, and the esophagus was mobilized off the crura and mediastinal structures. The anterior and posterior vagus nerves were identified and protected. Once 3 cm of intra-abdominal esophageal length was obtained, the



**Figure 2.** Operative imaging. (A) minimal adhesions and preservation of angle of His; (B) enhanced cardial wrap; (C) gastric retraction exposing left crus and posterior fundus; and (D) intact short gastric vessels.

crura were repaired primarily. The short gastric vessels were not divided (Figure 2D). The laparoscopic portion was concluded, and TIF was performed using the Esophyx device to create a 270-degree wrap (Figure 1C).

## Results

The operative time was 104 minutes. There were no intra-operative or postoperative complications. She was discharged the following day and discontinued PPI therapy 2 weeks postoperatively.

The patient presented 33 months later with upper abdominal pain and nausea. EGD showed no hiatal hernia and intact TIF wrap (Figure 1D). Further workup showed biliary dyskinesia. The patient underwent laparoscopic cholecystectomy which was uneventful. This provided the opportunity to examine the operative anatomy. There were minimal adhesions to the liver. The angle of His was preserved. The wrap appeared intact. The access to the left crus was straightforward.

The patient made an uneventful recovery. The epigastric pain and nausea resolved. The pathological examination showed chronic cholecystitis. The GERD-HRQL, RSI, and GERSS at 33 months off PPI were 0, 2, and 0, respectively.

## Discussion

LEHHR concept has many advantages. For simplicity, we list 10 reasons. They are categorized into 2 functional, 2 anatomical, 4 operative, and 2 strategic reasons.

First, LEHHR creates a partial wrap that is less likely to produce gas bloat syndrome or dysphagia. Second, this approach has been shown to be effective in normalizing esophageal pH scores.<sup>2</sup> Third, it preserves the angle of His which is an important antireflux mechanism. Fourth, it does not impact the blood supply of the stomach as it does not require division of the short gastric vessels. Fifth, it is less likely to cause bleeding from short gastric vessels or splenic tears. Sixth, it involves less operative dissection as it leaves the retrogastric plane undisturbed. Seventh, it provides an alternative when laparoscopic fundoplication is challenging. This was highlighted in recent reports of massive caudate lobe or prominent aberrant left hepatic artery.<sup>3,4</sup> LEHHR was beneficial in this patient with a small torso barely enough for 4 laparoscopic ports. Eighth, LEHHR produces a reproducible wrap which decreases variability among surgeons. A recent study evaluating a novel technique for the repair of the diaphragmatic hiatus utilized LEHHR to standardize the wrap thus focusing on the impact of cruroplasty.<sup>5</sup> Ninth, the Esophyx device wraps the cardia thus sparing the fundus for any potential subsequent antireflux surgery (Figures 1A and 2B). Tenth, the access to the left crus via

the retrogastric plane is a critical step in revisional antireflux surgery; this plane is undisturbed which facilitates possible future revisional surgery (Figure 2C).

## Conclusion

Laparoendoscopic hiatal hernia repair has functional, anatomical, operative, and strategic advantages. It provides a safe and effective alternative in challenging surgical scenarios. Prospective studies with longer follow-up are required to validate this technique.

## Acknowledgements

We are grateful to Dr. John Ferrara and Dr. Wei Wei for their insights regarding laparoendoscopic hiatal hernia repair. We thank Mrs. Sarah Williams and Mr. David Lorenson for their contribution in editing this manuscript.

## Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

## Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

## References

1. Ihde GM, Besancon K, Deljkich E. Short-term safety and symptomatic outcomes of transoral incisionless fundoplication with or without hiatal hernia repair in patients with chronic gastroesophageal reflux disease. *Am J Surg.* 2011;202(6):740-747. doi:10.1016/j.amjsurg.2011.06.035
2. Ihde GM, Pena C, Scitern C, Brewer S. pH scores in hiatal repair with transoral incisionless fundoplication. *JLS.* 2019;23(1):e2018.00087. doi:10.4293/JLS.2018.00087
3. Fanous M, Jaehne A, Lorenson D, Williams S. Massive caudate lobe in laparoscopic hiatal hernia repair: tying and wrapping outside the box. *Am Surg.* 2019;85(7):e336-e8.
4. Fanous MY. Benefit of laparoendoscopic repair of hiatal hernia in the presence of aberrant left hepatic artery. *JLS.* 2019;23(1):e2019.00004. doi:10.4293/JLS.2019.00004
5. Fanous M. Diaphragmatic crural eversion: mid-term data of a novel technique to optimize HiatoPlasty during laparoscopic repair of hiatal hernia. *Am Surg.* 2020;86(1):e17-e19.